

Oxygen therapy

View Online



[1]

Beurskens, C.J.P. et al. 2015. The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review. *Respiration*. 89, 2 (2015), 166–174. <https://doi.org/10.1159/000369472>.

[2]

BTS Emergency Oxygen Guideline Development Group 2017. BTS guideline for emergency oxygen use in adult patients. *Thorax*. 72, Supplement 1 (2017), i2–i89.

[3]

Cabello, J.B. et al. 2013. Oxygen therapy for acute myocardial infarction. *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd.

[4]

Curley, G.F. et al. 2015. Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation? *Journal of thoracic disease*. 7, 7 (2015).

[5]

Egi, M. et al. 2018. Oxygen management in mechanically ventilated patients: A multicenter prospective observational study. *Journal of Critical Care*. 46, (2018), 1–5. <https://doi.org/10.1016/j.jcrc.2018.03.024>.

[6]

Ferguson, N.D. 2016. Oxygen in the ICU. JAMA. (2016).
<https://doi.org/10.1001/jama.2016.13800>.

[7]

Girardis, M. et al. 2016. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit. JAMA. (2016).
<https://doi.org/10.1001/jama.2016.11993>.

[8]

Hernández, G. et al. 2016. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients. JAMA. (2016). <https://doi.org/10.1001/jama.2016.14194>.

[9]

Kallet, R.H. and Branson, R.D. 2016. Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients? Respiratory Care. 61, 6 (2016), 801–817.
<https://doi.org/10.4187/respcare.04933>.

[10]

Lee, J.H. et al. 2013. Use of high flow nasal cannula in critically ill infants, children, and adults: a critical review of the literature. Intensive Care Medicine. 39, 2 (2013), 247–257.
<https://doi.org/10.1007/s00134-012-2743-5>.

[11]

Page, D. et al. 2018. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. Critical Care. 22, 1 (2018).
<https://doi.org/10.1186/s13054-017-1926-4>.

[12]

Spoletini, G. et al. 2016. High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia. JAMA. 315, 13 (2016). <https://doi.org/10.1001/jama.2016.2709>.

[13]

Stéphan, F. et al. 2015. High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery. *JAMA*. 313, 23 (2015).
<https://doi.org/10.1001/jama.2015.5213>.

[14]

Stub, D. et al. 2015. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation*. 131, 24 (2015), 2143–2150.
<https://doi.org/10.1161/CIRCULATIONAHA.114.014494>.

[15]

Sztrymf, B. et al. 2012. Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study. *Journal of Critical Care*. 27, 3 (2012), 324.e9-324.e13. <https://doi.org/10.1016/j.jcrc.2011.07.075>.