

# Oxygen therapy

View Online



---

1

BTS Emergency Oxygen Guideline Development Group. BTS guideline for emergency oxygen use in adult patients. *Thorax*. 2017;72:i2-89.

2

Cabello JB, Burls A, Emparanza JI, et al. Oxygen therapy for acute myocardial infarction. *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd 2013.

3

Egi M, Kataoka J, Ito T, et al. Oxygen management in mechanically ventilated patients: A multicenter prospective observational study. *Journal of Critical Care*. 2018;46:1-5. doi: 10.1016/j.jcrc.2018.03.024

4

Page D, Ablordeppey E, Wessman BT, et al. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. *Critical Care*. 2018;22. doi: 10.1186/s13054-017-1926-4

5

Ferguson ND. Oxygen in the ICU. *JAMA*. Published Online First: 2016. doi: 10.1001/jama.2016.13800

6

Girardis M, Busani S, Damiani E, et al. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit. *JAMA*. Published Online First: 2016. doi: 10.1001/jama.2016.11993

7

Kallet RH, Branson RD. Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients? *Respiratory Care*. 2016;61:801–17. doi: 10.4187/respcare.04933

8

Stub D, Smith K, Bernard S, et al. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation*. 2015;131:2143–50. doi: 10.1161/CIRCULATIONAHA.114.014494

9

Hernández G, Vaquero C, Colinas L, et al. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients. *JAMA*. Published Online First: 2016. doi: 10.1001/jama.2016.14194

10

Spoletini G, Garpestad E, Hill NS. High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia. *JAMA*. 2016;315. doi: 10.1001/jama.2016.2709

11

Curley GF, Laffy JG, Zhang H, et al. Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation? *Journal of thoracic disease*. 2015;7.

12

Lee JH, Rehder KJ, Williford L, et al. Use of high flow nasal cannula in critically ill infants, children, and adults: a critical review of the literature. *Intensive Care Medicine*. 2013;39:247–57. doi: 10.1007/s00134-012-2743-5

13

Stéphan F, Barrucand B, Petit P, et al. High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery. *JAMA*. 2015;313. doi: 10.1001/jama.2015.5213

14

Sztrymf B, Messika J, Mayot T, et al. Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study. *Journal of Critical Care*. 2012;27:324.e9-324.e13. doi: 10.1016/j.jcrc.2011.07.075

15

Beurskens CJP, Wösten-van Asperen RM, Preckel B, et al. The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review. *Respiration*. 2015;89:166–74. doi: 10.1159/000369472