

Oxygen therapy

View Online



[1]

BTS Emergency Oxygen Guideline Development Group, 'BTS guideline for emergency oxygen use in adult patients', *Thorax*, vol. 72, no. Supplement 1, pp. i2–i89, 2017.

[2]

J. B. Cabello, A. Burls, J. I. Emparanza, S. Bayliss, and T. Quinn, 'Oxygen therapy for acute myocardial infarction', in *Cochrane Database of Systematic Reviews*, Chichester, UK: John Wiley & Sons, Ltd, 2013. doi: 10.1002/14651858.CD007160.pub3. Available: <http://doi.wiley.com/10.1002/14651858.CD007160.pub3>

[3]

M. Egi et al., 'Oxygen management in mechanically ventilated patients: A multicenter prospective observational study', *Journal of Critical Care*, vol. 46, pp. 1–5, 2018, doi: 10.1016/j.jcrc.2018.03.024

[4]

D. Page et al., 'Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study', *Critical Care*, vol. 22, no. 1, 2018, doi: 10.1186/s13054-017-1926-4

[5]

N. D. Ferguson, 'Oxygen in the ICU', *JAMA*, 2016, doi: 10.1001/jama.2016.13800

[6]

M. Girardis et al., 'Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit', *JAMA*, 2016, doi: 10.1001/jama.2016.11993

[7]

R. H. Kallet and R. D. Branson, 'Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients?', *Respiratory Care*, vol. 61, no. 6, pp. 801–817, 2016, doi: 10.4187/respcare.04933

[8]

D. Stub et al., 'Air versus oxygen in ST-segment-elevation myocardial infarction', *Circulation*, vol. 131, no. 24, pp. 2143–2150, 2015, doi: 10.1161/CIRCULATIONAHA.114.014494

[9]

G. Hernández et al., 'Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients', *JAMA*, 2016, doi: 10.1001/jama.2016.14194

[10]

G. Spoletini, E. Garpestad, and N. S. Hill, 'High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia', *JAMA*, vol. 315, no. 13, 2016, doi: 10.1001/jama.2016.2709

[11]

G. F. Curley, J. G. Laffy, H. Zhang, and A. S. Slutsky, 'Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation?', *Journal of thoracic disease*, vol. 7, no. 7, 2015.

[12]

J. H. Lee, K. J. Rehder, L. Williford, I. M. Cheifetz, and D. A. Turner, 'Use of high flow nasal cannula in critically ill infants, children, and adults: a critical review of the literature', *Intensive Care Medicine*, vol. 39, no. 2, pp. 247–257, 2013, doi: 10.1007/s00134-012-2743-5

[13]

F. Stéphan et al., 'High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery', *JAMA*, vol. 313, no. 23, 2015, doi: 10.1001/jama.2015.5213

[14]

B. Sztrymf, J. Messika, T. Mayot, H. Lenglet, D. Dreyfuss, and J.-D. Ricard, 'Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study', *Journal of Critical Care*, vol. 27, no. 3, p. 324.e9-324.e13, 2012, doi: 10.1016/j.jcrc.2011.07.075

[15]

C. J. P. Beurskens, R. M. Wösten-van Asperen, B. Preckel, and N. P. Juffermans, 'The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review', *Respiration*, vol. 89, no. 2, pp. 166–174, 2015, doi: 10.1159/000369472