

Oxygen therapy

View Online



1.

BTS Emergency Oxygen Guideline Development Group. BTS guideline for emergency oxygen use in adult patients. *Thorax*. 2017;72(Supplement 1):i2-i89.
2.

Cabello JB, Burls A, Emparanza JI, Bayliss S, Quinn T. Oxygen therapy for acute myocardial infarction. *Cochrane Database of Systematic Reviews* [Internet]. Chichester, UK: John Wiley & Sons, Ltd; 2013. Available from: <http://doi.wiley.com/10.1002/14651858.CD007160.pub3>
3.

Egi M, Kataoka J, Ito T, Nishida O, Yasuda H, Okamaoto H, Shimoyama A, Izawa M, Matsumoto S, Furushima N, Yamashita S, Takada K, Ohtsuka M, Fujisaki N, Shime N, Inagaki N, Taira Y, Yatabe T, Nitta K, Yokoyama T, Kushimoto S, Tokunaga K, Doi M, Masuda T, Miki Y, Matsuda K, Asaga T, Hazama K, Matsuyama H, Nishimura M, Mizobuchi S. Oxygen management in mechanically ventilated patients: A multicenter prospective observational study. *Journal of Critical Care*. 2018;46:1-5.
4.

Page D, Ablordeppey E, Wessman BT, Mohr NM, Trzeciak S, Kollef MH, Roberts BW, Fuller BM. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. *Critical Care*. 2018;22(1).
5.

Ferguson ND. Oxygen in the ICU. *JAMA*. 2016;

6.

Girardis M, Busani S, Damiani E, Donati A, Rinaldi L, Marudi A, Morelli A, Antonelli M, Singer M. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit. *JAMA*. 2016;

7.

Kallet RH, Branson RD. Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients? *Respiratory Care*. 2016;61(6):801–817.

8.

Stub D, Smith K, Bernard S, Nehme Z, Stephenson M, Bray JE, Cameron P, Barger B, Ellims AH, Taylor AJ, Meredith IT, Kaye DM. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation*. 2015;131(24):2143–2150.

9.

Hernández G, Vaquero C, Colinas L, Cuenca R, González P, Canabal A, Sanchez S, Rodriguez ML, Villasclaras A, Fernández R. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients. *JAMA*. 2016;

10.

Spoletini G, Garpestad E, Hill NS. High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia. *JAMA*. 2016;315(13).

11.

Curley GF, Laffy JG, Zhang H, Slutsky AS. Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation? *Journal of thoracic disease*. 2015;7(7).

12.

Lee JH, Rehder KJ, Williford L, Cheifetz IM, Turner DA. Use of high flow nasal cannula in

critically ill infants, children, and adults: a critical review of the literature. *Intensive Care Medicine*. 2013;39(2):247–257.

13.

Stéphan F, Barrucand B, Petit P, Rézaiguia-Delclaux S, Médard A, Delannoy B, Cosserant B, Flicoteaux G, Imbert A, Pilorge C, Bérard L. High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery. *JAMA*. 2015;313(23).

14.

Sztrymf B, Messika J, Mayot T, Lenglet H, Dreyfuss D, Ricard JD. Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study. *Journal of Critical Care*. 2012;27(3):324.e9-324.e13.

15.

Beurskens CJP, Wösten-van Asperen RM, Preckel B, Juffermans NP. The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review. *Respiration*. 2015;89(2):166–174.