

Oxygen therapy

View Online



-
1.
BTS Emergency Oxygen Guideline Development Group. BTS guideline for emergency oxygen use in adult patients. *Thorax* **72**, i2–i89 (2017).

 2.
Cabello, J. B., Burls, A., Emparanza, J. I., Bayliss, S. & Quinn, T. Oxygen therapy for acute myocardial infarction. in *Cochrane Database of Systematic Reviews* (John Wiley & Sons, Ltd, Chichester, UK, 2013). doi:10.1002/14651858.CD007160.pub3.

 3.
Egi, M. et al. Oxygen management in mechanically ventilated patients: A multicenter prospective observational study. *Journal of Critical Care* **46**, 1–5 (2018).

 4.
Page, D. et al. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. *Critical Care* **22**, (2018).

 5.
Ferguson, N. D. Oxygen in the ICU. *JAMA* (2016) doi:10.1001/jama.2016.13800.

 6.
Girardis, M. et al. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit. *JAMA* (2016) doi:10.1001/jama.2016.11993.

7.

Kallet, R. H. & Branson, R. D. Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients? *Respiratory Care* **61**, 801–817 (2016).

8.

Stub, D. et al. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation* **131**, 2143–2150 (2015).

9.

Hernández, G. et al. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients. *JAMA* (2016) doi:10.1001/jama.2016.14194.

10.

Spoletini, G., Garpestad, E. & Hill, N. S. High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia. *JAMA* **315**, (2016).

11.

Curley, G. F., Laffy, J. G., Zhang, H. & Slutsky, A. S. Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation? *Journal of thoracic disease* **7**, (2015).

12.

Lee, J. H., Rehder, K. J., Williford, L., Cheifetz, I. M. & Turner, D. A. Use of high flow nasal cannula in critically ill infants, children, and adults: a critical review of the literature. *Intensive Care Medicine* **39**, 247–257 (2013).

13.

Stéphan, F. et al. High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery. *JAMA* **313**, (2015).

14.

Sztrymf, B. et al. Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study. *Journal of Critical Care* **27**, 324.e9-324.e13 (2012).

15.

Beurskens, C. J. P., Wösten-van Asperen, R. M., Preckel, B. & Juffermans, N. P. The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review. *Respiration* **89**, 166-174 (2015).