

# Oxygen therapy

View Online



1.  
BTS Emergency Oxygen Guideline Development Group. BTS guideline for emergency oxygen use in adult patients. *Thorax*. 2017;72(Supplement 1):i2-89.
  
2.  
Cabello JB, Burls A, Emparanza JI, Bayliss S, Quinn T. Oxygen therapy for acute myocardial infarction. In: *Cochrane Database of Systematic Reviews* [Internet]. Chichester, UK: John Wiley & Sons, Ltd; 2013. Available from: <http://doi.wiley.com/10.1002/14651858.CD007160.pub3>
  
3.  
Egi M, Kataoka J, Ito T, Nishida O, Yasuda H, Okamaoto H, et al. Oxygen management in mechanically ventilated patients: A multicenter prospective observational study. *Journal of Critical Care*. 2018;46:1-5.
  
4.  
Page D, Ablordeppey E, Wessman BT, Mohr NM, Trzeciak S, Kollef MH, et al. Emergency department hyperoxia is associated with increased mortality in mechanically ventilated patients: a cohort study. *Critical Care*. 2018;22(1).
  
5.  
Ferguson ND. Oxygen in the ICU. *JAMA*. 2016;
  
- 6.

Girardis M, Busani S, Damiani E, Donati A, Rinaldi L, Marudi A, et al. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit. *JAMA*. 2016;

7.

Kallet RH, Branson RD. Should oxygen therapy be tightly regulated to minimize hyperoxia in critically ill patients? *Respiratory Care*. 2016;61(6):801–17.

8.

Stub D, Smith K, Bernard S, Nehme Z, Stephenson M, Bray JE, et al. Air versus oxygen in ST-segment-elevation myocardial infarction. *Circulation*. 2015;131(24):2143–50.

9.

Hernández G, Vaquero C, Colinas L, Cuenca R, González P, Canabal A, et al. Effect of Postextubation High-Flow Nasal Cannula vs Noninvasive Ventilation on Reintubation and Postextubation Respiratory Failure in High-Risk Patients. *JAMA*. 2016;

10.

Spoletini G, Garpestad E, Hill NS. High-flow nasal oxygen or noninvasive ventilation for postextubation hypoxemia. *JAMA*. 2016;315(13).

11.

Curley GF, Laffy JG, Zhang H, Slutsky AS. Noninvasive respiratory support for acute respiratory failure—high flow nasal cannula oxygen or non-invasive ventilation? *Journal of thoracic disease*. 2015;7(7).

12.

Lee JH, Rehder KJ, Williford L, Cheifetz IM, Turner DA. Use of high flow nasal cannula in critically ill infants, children, and adults: a critical review of the literature. *Intensive Care Medicine*. 2013;39(2):247–57.

13.

Stéphan F, Barrucand B, Petit P, Rézaiguia-Delclaux S, Médard A, Delannoy B, et al. High-flow nasal oxygen vs noninvasive positive airway pressure in hypoxemic patients after cardiothoracic surgery. *JAMA*. 2015;313(23).

14.

Sztrymf B, Messika J, Mayot T, Lenglet H, Dreyfuss D, Ricard JD. Impact of high-flow nasal cannula oxygen therapy on intensive care unit patients with acute respiratory failure: A prospective observational study. *Journal of Critical Care*. 2012;27(3):324.e9-324.e13.

15.

Beurskens CJP, Wösten-van Asperen RM, Preckel B, Juffermans NP. The potential of heliox as a therapy for acute respiratory distress syndrome in adults and children: a descriptive review. *Respiration*. 2015;89(2):166-74.